

██████████
████████████████████
██

Certificate of Mold Analysis

Prepared for: TEXINSPEC
Phone Number:
Fax Number:
Project Name: ██████████
Test Location: ██████████
██
Chain of Custody #: ██████████
Received Date: November 6, 2018
Report Date: November 6, 2018

Elena Santiago, Technical Manager

Olmedo Ferrer, Quality Control Manager

Currently there are no Federal regulations for evaluating potential health effects of fungal contamination and remediation. This information is subject to change as more information regarding fungal contaminants becomes available. For more information visit <http://www.epa.gov/mold> or www.nyc.gov/html/doh/html/epi/mold.shtml. This document was designed to follow currently known industry guidelines for the interpretation of microbial sampling, analysis, and remediation. Since interpretation of mold analysis reports is a scientific work in progress, it may as such be changed at any time without notice. The client is solely responsible for the use or interpretation. PRO-LAB/SSPTM Inc. makes no express or implied warranties as to health of a property from only the samples sent to their laboratory for analysis. The Client is hereby notified that due to the subjective nature of fungal analysis and the mold growth process, laboratory samples can and do change over time relative to the originally sampled material. PRO-LAB/SSPTM Inc. reserves the right to properly dispose of all samples after the testing of such samples are sufficiently completed or after a 7 day period, whichever is greater.



Lab ID # 163230

For more information please contact PRO-LAB at (954) 384-4446 or email info@prolabinc.com

Prepared for : TEXINSPEC

Test Address : [REDACTED]
[REDACTED]
[REDACTED]

ANALYSIS METHOD	6110 Air Direct Examination	6110 Air Direct Examination	6110 Air Direct Examination	INTENTIONALLY BLANK
LOCATION	OUTSIDE	KITCHEN	REAR BEDROOM	
COC / LINE #	1184768-1	1184768-2	1184768-3	
SAMPLE TYPE & VOLUME	AIR-O-CELL - 150L	AIR-O-CELL - 150L	AIR-O-CELL - 150L	
SERIAL NUMBER	26806240	26806249	26806273	
COLLECTION DATE	Nov 5, 2018	Nov 5, 2018	Nov 5, 2018	
ANALYSIS DATE	Nov 6, 2018	Nov 6, 2018	Nov 6, 2018	
CONCLUSION	CONTROL	NOT ELEVATED	ELEVATED	

IDENTIFICATION	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total
Alternaria				8	53	5						
Bipolaris/Drechslera	4	27	2									
Cladosporium	64	430	24	28	190	19	28	190	15			
Nigrospora	4	27	2									
Other Ascospores	40	270	15									
Other Basidiospores	112	750	42									
Penicillium/Aspergillus				116	770	76	164	1,100	85			
Peronospora	4	27	2									
Rusts	8	53	3									
Smuts, myxomycetes	32	210	12									

TOTAL SPORES	268	1,794	100	152	1,013	100	192	1,290	100			
MINIMUM DETECTION LIMIT*	4	27		4	27		4	27				

BACKGROUND DEBRIS	Light			Light			Light					
Cellulose Fiber	4	27		4	27		4	27				
Pollen	4	27										

OBSERVATIONS & COMMENTS				

Background debris qualitatively estimates the amount of particles that are not pollen or spores and directly affects the accuracy of the spore counts. The categories of Light, Moderate, Heavy and Too Heavy for Accurate Count, are used to indicate the amount of deposited debris. Light (None to up to 25% obstruction); Medium (26% to up to 75% obstruction); Heavy (76% to up to 90% obstruction); Too Heavy (Greater than 90% obstruction). Increasing amounts of debris will obscure small spores and can prevent spores from impacting onto the slide. The actual number of spores present in the sample is likely higher than reported if the debris estimate is 'Heavy' or 'Too Heavy for Accurate Count'. All calculations are rounded to two significant figures and therefore, the total percentage of spore numbers may not equal 100%. The effect of the results relate only to the items tested. The methods used in this analysis have been validated and is fit for the intended use. R "version" indicated after the lab D# indicates a sample with amended data.

* Minimum Detection Limit. Based on the volume of air sampled, this is the lowest number of spores that can be detected and is an estimate of the lowest concentration of spores that can be read in the sample. NA = Not Applicable.

Spores that were observed from the samples submitted are listed on this report. If a spore is not listed on this report it was not observed in the samples submitted.

Interpretation Guidelines A determination is added to the report to help users interpret the mold analysis results. A mold report is only one aspect of an indoor air quality investigation. The most important aspect of mold growth in a living space is the availability of water. Without a source of water, mold generally will not become a problem in buildings. These determinations are in no way meant to imply any health outcomes or financial decisions based solely on this report. For questions relating to medical conditions you should consult an occupational or environmental health physician or professional.

CONTROL is a baseline sample showing what the spore count and diversity is at the time of sampling. The control sample(s) is usually collected outside of the structure being tested and used to determine if this sample(s) is similar in diversity and abundance to the inside sample(s).


ELEVATED means that the amount and/or diversity of spores, as compared to the control sample(s), and other samples in our database, are higher than expected. This can indicate that fungi have grown because of a water leak or water intrusion. Fungi that are considered to be indicators of water damage include, but are not limited to: *Chaetomium*, *Fusarium*, *Memnoniella*, *Stachybotrys*, *Scopulariopsis*, *Ulocladium*.

NOT ELEVATED means that the amount and/or the diversity of spores, as compared to the control sample and other samples in our database, are lower than expected and may indicate no problematic fungal growth.

UNUSUAL means that the presence of current or former growth was observed in the analyzed sample. An abundance of spores are present, and/or growth structures including hyphae and/or fruiting bodies are present and associated with one or more of the types of mold/fungi identified in the analyzed sample.

Chain of Custody # [Redacted]


[Green checkered pattern] Kitchen
[Blue solid] Outside



Chain of Custody # [Redacted]

[Green Checkered Box] Rear Bedroom

[Blue Box] Outside



Identification	Outdoor Habitat	Indoor Habitat	Possible Allergic Potential Not an opinion or interpretation	Comments
Alternaria	One of the most commonly reported airborne spores worldwide. Often common in outdoor air. Usually not observed in large numbers in outdoor air. Soil, dead or dying plants, foodstuffs, textiles	Wallboard paper backing, wood, other various cellulose-containing materials. Commonly found in settled dust and as normal settled spores on carpets, drapes, textiles, etc.	Common allergen. Type I allergies (hay fever and asthma); Type III hypersensitivity pneumonitis. Common cause of extrinsic asthma.	Alternaria is commonly found in elevated numbers on water-intruded building materials and in higher spore numbers in the air with respect to the outside when growth on wet building materials occurs.
Bipolaris/Drechslera	Common everywhere. Frequently associated with grasses, but also found on plant material, decaying food, and soil.		Common Type I (hay fever and asthma), fungal sinusitis.	This is a group of like-looking spores that include Bipolaris, Drechslera, Exserohilum, and sometimes Helminosporium. They cannot be consistently separated by spore morphology and are thus grouped together. Must be cultured to consistently separate the genera.
Cladosporium	The most common spore type reported in the air worldwide. Found on dead and dying plant litter, and soil.	Commonly found on wood and wallboard. Commonly grows on window sills, textiles and foods.	Type I (hay fever and asthma), Type III (hypersensitivity pneumonitis) allergies.	A very common and important allergen source both outdoors and indoors.
Nigrospora	Commonly found everywhere. Grows on decaying plant material	Does not normally grow on building materials, but occasionally can be found growing on wallboard.	Type I (hay fever and asthma) allergies.	Very distinctive spore that is easy to identify.
Ascospores	Common everywhere. Constitutes a large part of the airspora outside. Can reach very high numbers in the air outside during the spring and summer. Can increase in numbers during and after rainfalls.	Very few of this group grow inside. The notable exception is Chaetomium, Ascotricha and Peziza.	Little known for most of this group of fungi. Dependent on the type (see Chaetomium and Ascotricha).	
Basidiospores	Commonly found everywhere, especially in the late summer and fall. These spores are from Mushrooms.	Mushrooms are not normally found growing indoors, but can grow on wet lumber, especially in crawlspaces. Sometimes mushrooms can be seen growing in flower pots indoors.	Some allergenicity reported. Type I (hay fever, asthma) and Type III (hypersensitivity pneumonitis).	Among the group of Mushrooms (Basidiomycetes) are dry rot fungi Serpula and Poria that are particularly destructive to buildings.
Penicillium/Aspergillus	Common everywhere. Normally found in the air in small amounts in outdoor air. Grows on nearly everything.	Wetted wallboard, wood, food, leather, etc. Able to grow on many substrates indoors.	Type I (hay fever and asthma) allergies and Type III (hypersensitivity pneumonitis) allergies.	This is a combination group of Penicillium and Aspergillus and is used when only the spores are seen. The spores are so similar that they cannot be reliably separated into their respective genera.
Peronospora	Rarely seen in the air. Grows on leave, stems and fruits of living plants.	None known.	None known.	Needs a living host to grow.

Identification	Outdoor Habitat	Indoor Habitat	Possible Allergic Potential Not an opinion or interpretation	Comments
Rusts	Common everywhere growing on grasses, trees and other living plants.	Does not grow indoors.	Type I (hay fever and asthma) allergies.	Rust requires a living plant host to complete part of its lifecycle and thus, is not normally found growing indoors except perhaps on an infected house plant.
Smuts, myxomycetes	Commonly found everywhere, especially on logs, grasses and weeds.	Smuts don't normally grow indoors, but can occasionally be found on things brought from outside and stored in the house. Myxomycetes can occasionally grow indoors, but need lots of water to be established.	Type I (hay fever and asthma) allergies.	Smuts and myxomycetes are a combined group of organisms because their spores look so similar and cannot be reliably distinguished from each other.